

10:20 am ~

多相星間媒質における 巨大分子雲質量関数の時間発展と 分子雲衝突に誘起された星形成

(Kobayashi+ 2017 in ApJ, 2018 in revision in PASJ, 2018 in progress.)

Masato I.N. KOBAYASHI
(D3, Nagoya U)



Team BISTRO-J

Rironkon Symposium 2017, Dec. 26th 2017 at U of Tokyo

Collaborators; Shu-ichiro Inutsuka, Hiroshi Kobayashi,
Yasuo Fukui, Kenji Hasegawa,
Tsuyoshi Inoue (Nagoya U)

Outline

- ✓ **Background**
 - ◆ **Multiple Episodes of Compression** in magnetized ISM

- ✓ **Time Evolution of GMCs on Galactic Scales**
 - ◆ **Network of expanding shells**
 - ◆ Formulation of GMC MF evolution with Cloud-Cloud Collisions (CCC)

- ✓ **CCC-driven (Massive) Star Formation**
 - ◆ **Rapid dispersal of GMCs**

- ✓ **Towards Galactic-Scale Studies**
 - ◆ Shock propagation in the multiphase ISM

- ✓ **Summary**


Backgrounds

- ✓ **Observed GMC MF**
- ✓ **Multiphase Simulations**
- ✓ **Multiple Episodes of Compressions**


IRAM: GMC Distribution in M51

Sub-division in the galactic disk (Colombo+ 2014a)

1) Face-on view of identified GMCs



2) Cumulative mass function of GMCs




Power-law slope variation:
 shallower (< -2) in **arm** regions,
 steeper (> -2) in **inter-arm** regions



➡ Larger GMCs tend to reside along spirals.
 (Consistent trend from Koda+ 2009)

Magnetic fields retard cloud formation

Inoue & Inutsuka, 2008



Inoue & Inutsuka, 2012




Cloud formation take place after supersonic **compression with multiple times.**

ISM Simulations

Typical time scale for GMC formation

Multiple episodes of compression is essential to form molecular clouds from magnetized WNM, which occupy most of the volume in galactic disks.



Goal: determine how this multiple compression governs the observed variation in GMC mass functions.

Time Evolution of GMCs on Galactic Scales


- ✓ **Network of expanding shells**
- ✓ **Formulation of GMC MF evolution**

GMC Formation & Evolution

Network of expanding shells on galactic scales

(Inutsuka+ 2015 A&A)

HII region/ SNR



Cloud-Cloud Collisions (CCCs)



Fukui+ 2014
Nakamura+ 2012
Torii+ 2015

➔ Formulate the **time evolution of GMC populations** coarse-grained on a few 10 to **100 pc scales** (disk regions).

Formulation

(c.f., Levinson & Roberts 1981, Kwan 1979, Scoville & Hersch 1979, Cowie 1980, Tomisaka 1984)



n_{cl} : differential number density of GMCs with mass m

$$\frac{\partial n_{cl}}{\partial t} + \frac{\partial}{\partial m} \left(n_{cl} \frac{m}{T_f} \right) =$$

$$-\frac{n_{cl}}{T_d}$$

GMC self-dispersal

due to radiation by massive stars
($T_d \sim 14$ Myr; c.f. radiation hydrodynamics simulation: Inutsuka 2015)

$$+ \frac{1}{2} \int_0^\infty \int_0^\infty K(m_1, m_2) n_{cl,1} n_{cl,2} \times \delta(m - m_1 - m_2) dm_1 dm_2$$

$$- \int_0^\infty K(m, m_2) n_{cl} n_{cl} dm_2$$

$$+ \frac{1}{m} \frac{\partial (n_{cl} m)}{\partial t} \Big|_{res}$$

Gas Resurrection


(Replenishment of minimum-mass population)

GMC formation/growth
through compressions


($T_f \sim 10$ Myr;
c.f. Inoue & Inutsuka 2012)



Cloud-Cloud Collisions (CCCs)
(coagulation)




Dispersed Compressions



Fate of Dispersed Gas

Dispersed gas may resurrect to replenish the minimum-mass population.



Total gas dispersal rate
due to stellar feedback

Minimum-mass
in the system

$$\left. \frac{\partial (n_{c1}m)}{\partial t} \right|_{\text{res}} = \underbrace{\varepsilon_{\text{res}}}_{\text{↑}} \underbrace{\dot{\rho}_{\text{total,disp}}}_{\text{↓}} \delta(m - \underbrace{m_{\text{min}}}_{\text{↓}})$$

Resurrecting factor (0.01-1)

The rate to generate new generation minimum-mass GMCs out of the total dispersed gas.

N.B.: In a steady state, $1 - \varepsilon_{\text{res}}$ can be understood as a rate to accrete onto pre-existing GMCs out of the total dispersed gas.

Results

- ✓ **Slope of GMC MF**
- ✓ **Gas Resurrection**


Steady State with various T_f

CCC affects only the massive-end evolution...



$$\frac{\partial n_{\text{cl}}}{\partial t} + \frac{\partial}{\partial m} \left(n_{\text{cl}} \frac{m}{T_f} \right) = -\frac{n_{\text{cl}}}{T_d} \quad \Rightarrow \quad n_{\text{cl}}(m) = n_0 \left(\frac{m}{M_\odot} \right)^{-1 - \frac{T_f}{T_d}}$$

Arm: $T_f = 4.2$ Myr, $\varepsilon_{\text{res}} = 0.012$, Inter-arm: $T_f = 22.4$ Myr, $\varepsilon_{\text{res}} = 0.45$

Kobayashi+ 2017





Kobayashi+ 2017



Arm regions: **short T_f** due to many massive stars/supernovae, **little resurrection** due to many massive (i.e. large) GMCs.

Optimal Steady State Resurrection


Relation between resurrection and GMCMF slope



Large surveys may put unique **constraints on GMC formation/dispersal timescales and the resurrecting factor by measuring GMCMF slopes.**

Optimal Steady State Resurrection


Relation between resurrection and **GMCMF** slope



N.B.

(We suggest what resurrecting factor properly reproduces observations.)


Resurrecting processes of **optically thick H I** and **CO-dark H₂** should be confirmed/understood **in the context of physics** by massive simulations.



Large surveys may put unique **constraints on GMC formation/dispersal timescales** and the resurrecting factor by measuring **GMCMF slopes**.

Controversy on GMC lifetime

GMC self-growth time starting with 100 Msun



➔ Massive GMCs are long-lived;
Typical “Age” and typical “Lifetime” is different!

CCC-driven Star Formation




- ✓ **Rapid Massive Star Formation**
- ✓ **Star Formation Rate**
- ✓ **Collision Frequency**

Star Formation at CCC sites

O(B) stars and YSOs are often observed.

Ex.) RCW38 (Fukui+ 2016)

O(B) star candidates form at the intersection.




**1) Complementary distribution on the sky and
2) bridging feature in CO position-velocity diagram
indicate a recent CCC event ~ 0.1 Myr ago.**

Star Formation at CCC sites


O(B) star formation possibly triggered by CCC.

Ex.) **NGC3603** Similar trends observed in different sites (Westerlund2, W51, M17, NGC6334, NGC6357, M16, W33, M42, RCW166, S116, S117, S118, M43, RCW36, M20, RCW120, NGC2024, RCW34, ... etc.)

12CO contour + JHK image
Fukui+ 2014



Probability of stellar ages
(with assumed isochrones).




Multiple O stars might form **~0.5pc scale within 1Myr!**

➡ **Is this process important for galactic star formation...?**


Two GMC Populations

Short Td for GMCs undergoing CCC

1) Normal Population



2) CCC Population



→ Growth through HI accretion
 → CCC
 → Dispersal due to massive stars

- ✓ GMCs join the CCC population once they experience CCC.
- ✓ CCC population has a short Td $\sim 5\text{Myr}$
representing observed rapid massive-star formation.

Two GMC Populations

Time evolution equation and star formation rate (SFR)

n_{acc} : normal population

n_{col} : CCC population

$T_d = 14 \text{ Myr}$

$T_{d,\text{col}} = 5 \text{ Myr}$

$$\begin{aligned}
 & \frac{\partial (n_{\text{acc,cl}} + n_{\text{col,cl}})}{\partial t} + \frac{\partial}{\partial m} \left((n_{\text{acc,cl}} + n_{\text{col,cl}}) \frac{m}{T_f} \right) \\
 = & -\frac{n_{\text{acc,cl}}}{T_d} - \frac{n_{\text{col,cl}}}{T_{d,\text{col}}} \\
 & + \frac{1}{2} \int_0^\infty \int_0^\infty K(m_1, m_2) \\
 & \quad \times (n_{\text{acc,cl,1}} + n_{\text{col,cl,1}})(n_{\text{acc,cl,2}} + n_{\text{col,cl,2}}) \\
 & \quad \times \delta(m - m_1 - m_2) dm_1 dm_2 \\
 & - \int_0^\infty K(m, m_2) \\
 & \quad \times (n_{\text{acc,cl}} + n_{\text{col,cl}})(n_{\text{col,cl,2}} + n_{\text{col,cl,2}}) dm_2 \\
 & + \frac{1}{m} \frac{\partial (n_{\text{cl}} m)}{\partial t} \Big|_{\text{res}} .
 \end{aligned}$$


$\epsilon_{\text{SFE}} = 1\%$

$$\text{SFR}(> m) = \epsilon_{\text{SFE}} \times \left(\int_m^\infty \frac{m n_{\text{acc,cl}}}{T_d} dm + \int_m^\infty \frac{m n_{\text{col,cl}}}{T_{d,\text{col}}} dm \right)$$

CCC-driven star formation

Two GMC Populations

GMC MF and CCC-driven star formation rate



10kpc x 10kpc x 100pc disk (like Milky Way), this SFR = a few Msun per year.

→ **CCC-driven star formation may amount to a few 10 percent of the total star formation in the Milky Way and nearby galaxies, which is mostly driven by GMCs $> 10^5$ Msun.**

Which mass pair can we observe?

Observability


$$N_{\text{obs}}(m, m_2) = \int \frac{\int_m^{m+\delta m} n_{\text{cl}} dm}{T_{\text{col,num}}(m, m_2)} \Delta T_{\text{CCC}} dV_{\text{survey}}$$

ΔT_{CCC} : Duration over which CCC can be identified as CCC

$V_{\text{survey}} = \int dV_{\text{survey}}$: Survey volume

Let us optimistically assume $\Delta T_{\text{CCC}} = 1\text{Myr}$, and survey the entire MW disk.

Kobayashi+ 2018 in revision



➔ Most of the CCC pairs that **current observations** can probe must **consist of GMCs $\sim 10^4$ Msun**; we need to push **observations further to the pairs $> 10^5$ Msun**, which are important for galactic star formation!

Summary

✓ Backgrounds: **Multiple Episodes of Compression**

- ◆ Variation of GMC MF on galactic scales
- ◆ Magnetic fields retard molecular cloud formation

✓ Formulation: **Coagulation Equation with CCC**

- ◆ **GMC MF slope is characterized by T_f/T_d whereas its massive end is governed by CCC**
- ◆ **CCC-driven SF may amount to a few 10 per cent of total SF in the Milky Way galaxy.**

✓ Future Prospects:

- ◆ The column density of star formation rate (SFR)
- ◆ Transition from arm regions to inter-arm regions

