


Beyond a pale blue dot

Yasushi Suto

*Department of Physics and Research Center
for the Early Universe, the University of Tokyo*

November 12, 2015: The 2015 Kyoto Prize Workshop in Basic Sciences
"Formation mechanisms of planetary systems and the quest for earth-twins"



Revolution of our world-view

"We didn't know anything"



"Invincible"

Photo taken by Y.Suto
(Nov.4, 2004)

post-Nishinomiya-Yukawa
symposium, Kyoto University

Nightfall: We didn't know anything


A short novel by Issac Asimov



(Illustration: Alisa Haba)

- No “night” except the total eclipse due to an inner planet every 2049 years on the planet “Lagash”
- People realized the true world for the first time

History of exoplanet discovery



As of November 11, 2015 <http://exoplanet.eu/>

Are we alone ?

*a Pale Blue Dot ?
or pale blue dots ?*





Sciences with exoplanets

- the **final** question: *Are we alone ?*
 - origin of the earth
 - origin of the Solar System
 - **habitable** planets \Rightarrow origin of life
 - signature of **extra-terrestrial life** ?
 - \Rightarrow extra-terrestrial intelligence ?

“Where are they ?” E.Fermi (1950)

Solar planets imaged by Voyager 1 (February 4, 1990)


- Earth imaged at distance of 40 AU away
- A Pale Blue Dot coined by Carl Segan



A Pale Blue Dot



Voyager 1



Earth and Moon from Saturn (2013)



- Viewed from *Cassini* on July 20, 2013
 - about 20,000 happy Americans are waving their hand towards Cassini, but *how can we know that?*

Can we detect signatures
of life on our Earth ?


Search for signatures of life on "Earth" with Galileo mission! (1990)

- Launched in May, 1986
- Earth observed on December 8, 1990
- ***Conclusion: it is likely that life exists on Earth !***
 - Abundant O₂
 - Red-edge of vegetation
 - CH₄ abundance out of thermal equilibrium
 - Artificial pulsed radio signal



Sagan, Thompson,
Carlson, Gurnett & Hord:
Nature 365(1993)715

Conventional bio-signatures




- O₂
 - A-band@0.76μm
 - B-band@0.69μm
- H₂O
 - 0.72, 0.82, 0.94μm
- O₃
 - Chappuis band @ (0.5-0.7)μm
 - Hartley band @ (0.2-0.3)μm

Kasting et al. arXiv:0911.2936


“Exoplanet characterization and the search for life”

Sagan et al. (1993): spectrum of atmosphere



Strong O₂ absorption @A-band(0.76μm)


FIG. 1 a, Galileo long-wavelength-visible and near-infrared spectra of the Earth over a relatively cloud-free region of the Pacific Ocean, north of Borneo. The incidence and emission angles are 77° and 57° respectively. The $(b' \sum_g^+ - X^3 \sum_g^-)$ O-O band of O₂ at $0.76 \mu\text{m}$ is evident, along with a number of H₂O features. Using several cloud-free regions of varying airmass, we estimate an O₂ vertical column density of $1.5 \text{ km-atmag} \pm 25\%$. b and c, Infrared spectra of the Earth in the $2.4\text{--}5.2 \mu\text{m}$ region. The strong ν_3 CO₂ band is seen at the $4.3 \mu\text{m}$, and water vapour bands are found, but not indicated, in the $3.0 \mu\text{m}$ region. The ν_3 band of nitrous oxide, N₂O, is apparent at the edge of the CO₂ band near $4.5 \mu\text{m}$, and N₂O combination bands are also seen near $4.0 \mu\text{m}$. The



methane (0010) vibrational transition is evident at $3.31 \mu\text{m}$. A crude estimate¹⁰ of the CH₄ and N₂O column abundances is, for both species, of the order of 1 cm-atmag ($\equiv 1 \text{ cm path at STP}$).

Red edge of **(exo)plants**: a possible biosignature in **exoplanets**

- **Red-edge**
 - Significant increase of reflectivity of leaves on Earth (terrestrial planets) for $\lambda > 7000\text{\AA}$
- Widely used in the remote-sensing of our Earth



Seager, Ford & Turner
astro-ph/0210277

Vesto Melvin Slipher (1875-1969)



- Discovered redshifts of galaxies and thus cosmic expansion via the Doppler method

**“Observations of Mars in 1924 made at the Lowell Observatory: II spectrum observations of Mars’ ’
PASP 36(1924)261**




reflection spectrum. The Martian spectra of the dark regions so far do not give any certain evidence of the typical reflection spectrum of chlorophyl. The amount and types of vegetation required to make the effect noticeable is being investigated by suitable terrestrial exposures. ***Astrobiology indeed in 1924 !***

Sagan et al. (1993): colors of the earth

**Red-edge of the vegetation on the earth detected
by the Galileo mission**




Reflectivity



Sagan et al. (1993): radio observation


Detection of pulsed radio signals unlikely of natural/astronomical origin

FIG. 4 A frequency–time spectrogram of the radio signals detected by the Galileo plasma wave instrument. The intensities are coded in the sequence blue–green–yellow–red, with blue lowest and red highest. Several natural sources of radio emission are shown in *a*, including auroral kilometric radiation (AKR). Modulated emission at $f > 4$ MHz is shown with an expanded time scale in *b*. Modulated patterns of this type are characteristic of the transmission of information, and would be highly unusual for a naturally occurring radio source. (UT, universal time; R is distance of Galileo from Earth in units of Earth's radius, R_E ; LT, local time.)



Arecibo message (1974)


- Frank Drake sent a **radio message from Arecibo radio observatory** on November 16, 1974 towards globular cluster **M13 (25,000 light-year away)**
- The message, if decoded properly, should look like this.



- 1 to 10 in binary
- Atomic numbers of H, C, N, O, P that form DNA in binary
- Formulas for the sugars and bases in the nucleotides of DNA
- Double helix of DNA
- Human and the human population on earth
- The solar system
- Arecibo radio telescope

Simulated Earth
observed at 10pc away

Habitable zone around host stars



Kasting, Kopparapu, Ramirez & Harman: arXiv:1312.1328


Occurrence of earth-size habitable planets around Sun-like stars

- Planets with (1-2) Earth radius around GKstars
 - Kepler Transit planets corrected for selection effect
 - 11 ± 4 % (1-4 times the Solar flux on the earth)
 - $5.7 + 2.2 - 1.7$ % (orbital period of 200-400days)

Table 1. Occurrence of small planets in the habitable zone


HZ definition	a_{inner}	a_{outer}	$F_{P,\text{inner}}$	$F_{P,\text{outer}}$	f_{HZ} (%)
Simple	0.5	2	4	0.25	22
Kasting (1993)	0.95	1.37	1.11	0.53	5.8
Kopparapu et al. (2013)	0.99	1.70	1.02	0.35	8.6
Zsom et al. (2013)	0.38		6.92		26*
Pierrehumbert and Gaidos (2011)		10		0.01	$\sim 50^{\dagger}$

Starshade project: direct imaging of a second earth



Space telescope + occulting satellite at 50,000km away!
(Princeton Univ. + JPL/Caltech)

Expected daily change of the reflected light of the earth



Ford, Seager & Turner: Nature 412 (2001) 885

- Assume that the earth's reflected light is completely separated from the Sun's flux !
- Periodic change of 10% level due to different reflectivity of land, ocean, forest
- Cloud is the most uncertain factor: weather forecast !

Colors of a Second Earth: estimating the fractional areas of ocean, land and vegetation of Earth-like exoplanets

ApJ. 715(2010)866, arXiv:0911.5621

Colors of a Second Earth. II: Effects of Clouds on Photometric Characterization of Earth-like Exoplanets


ApJ. 738(2011)184, arXiv:1102.3625

- **Yuka Fujii**, H.Kawahara, A.Taruya, Y.Suto (Dept. of Phys., Univ. of Tokyo), S.Fukuda, T.Nakajima (Univ. of Tokyo, Center of climate system research), Edwin Turner (Princeton Univ.)


<http://www.space.com/scienceastronomy/color-changing-planets-alien-life-100513.html>

Colors of a second earth


- **Beyond a pale blue dot**
 - Impossible to spatially resolve the surface of a second earth
 - Color should change due to the rotation
 - A second earth = a **color-changing** dot



Colors of our earth




March 18th-19th






A pale blue dot ? Not really




Simulated
photometric
light-curves
of Earth

- Adopted Earth data in March
- Spin inclination = 0 (vernal equinox)
- cloudless




Fujii et al. (2010)

Estimating fractional areas of surface components from colors of a second earth



- 2 week observation of a cloudless Earth at 10 pc away
 - Reasonably well reproduced
 - possible to identify vegetation !
- Fujii et al. (2010)**


Surface latitude map estimated from real satellite data with cloud model



Old
M-star



Young
M-star



G-star



F-star



"The color of plants on other worlds"
N.Kiang, Scientific American (2008)

Summary:

A pale blue dot? Not really !

- Future direct imaging of **daily change of colors of another earth is challenging, but** would reveal the presence of ocean, land, cloud, and/or even vegetation on their surface
- **Detection of a second Earth may not be a mere fairy tale nor a science fiction any more**
- Detection of oxygen, water vapor, and even the red-edge may be a promising path towards astrobiology